

COMPACTLY-ALIGNED DISCRETE PRODUCT SYSTEMS, AND GENERALIZATIONS OF $\mathcal{O}_\infty$

NEAL J. FOWLER

ABSTRACT. The universal C^* -algebras of discrete product systems generalize the Toeplitz-Cuntz algebras and the Toeplitz algebras of discrete semigroups. We consider a semigroup P which is quasi-lattice ordered in the sense of Nica, and, for a product system $p : E \rightarrow P$, we study those representations of E , called covariant, which respect the lattice structure of P . We identify a class of product systems, which we call compactly aligned, for which there is a purely C^* -algebraic characterization of covariance, and study the algebra $C_{\text{cov}}^*(P, E)$ which is universal for covariant representations of E . Our main theorem is a characterization of the faithful representations of $C_{\text{cov}}^*(P, E)$ when P is the positive cone of a free product of totally-ordered amenable groups.

INTRODUCTION

The study of C^* -algebras generated by isometries has a rich history, beginning with Coburn's theorem that any two algebras $C^*(U)$ and $C^*(V)$ generated by nonunitary isometries U and V are canonically isomorphic [3]. Cuntz generalized Coburn's result by studying the C^* -algebra generated by a *Toeplitz-Cuntz* family $\{S_1, \dots, S_n\}$ of isometries with mutually orthogonal ranges; this led to his well-known analysis of the Cuntz algebra $\mathcal{O}_n$ and its Toeplitz extension $\mathcal{TO}_n$ [4, 5].

One can also view Coburn's algebra as the Toeplitz algebra of $\mathbb{N}$, and it is natural to seek versions of his theorem for other semigroups. A modern approach is to take a semigroup P , construct a C^* -algebra $C^*(P)$ which is universal for representations $V : P \rightarrow \text{Isom}(\mathcal{H})$ (called *isometric representations*), and ask when the integrated form $V_* : C^*(P) \rightarrow \mathcal{B}(\mathcal{H})$ is faithful; in this language Coburn's theorem says that $V_* : C^*(\mathbb{N}) \rightarrow \mathcal{B}(\mathcal{H})$ is faithful if and only if none of the isometries V_n is unitary. Douglas generalized this result by showing that one can replace $\mathbb{N}$ with the positive cone Γ^+ of any countable subgroup Γ of $\mathbb{R}$ [8], and Murphy later extended this to handle any totally-ordered abelian group Γ [12].

Here we combine the approaches of Cuntz and Douglas/Murphy by studying the C^* -algebra generated by an entire collection of Toeplitz-Cuntz families, one for each element of a semigroup P . The algebraic relations among the families are governed by a *product system* E over P , which, loosely speaking, is a collection $\{E_s : s \in P\}$ of Hilbert spaces, together with an associative multiplication which induces unitary isomorphisms $E_s \otimes E_t \rightarrow E_{st}$. (Our E is a discrete analogue of the continuous tensor product systems which arose in Arveson's work on one-parameter semigroups of endomorphisms of

Date: September 26, 1998.

1991 Mathematics Subject Classification. Primary 46L55.

$\mathcal{B}(\mathcal{H})$ [2].) Every representation of E gives a collection of Toeplitz-Cuntz families indexed by P , and with an appropriate choice of E we can realize any of the above Toeplitz algebras as the C^* -algebra $C^*(P, E)$ which is universal for representations of E : when E is the trivial product system over P we have $C^*(P, E) = C^*(P)$, and by taking the n -dimensional product system over $\mathbb{N}$ we have $C^*(P, E) = \mathcal{TO}_n$. (See [9, §1] for these and other examples.)

The C^* -algebra of a discrete product system was first studied by Dinh, who established the simplicity of $C^*(\Gamma^+, E)$ for any product system E over the positive cone Γ^+ of a countable dense subgroup Γ of $\mathbb{R}$, so long as each fiber E_s is infinite dimensional [6]. In [9], Fowler and Raeburn considered product systems over more general semigroups, but focussed primarily on systems with finite-dimensional fibers. Here we emphasize product systems which have infinite-dimensional fibers, and one of our goals is to extend Dihn's theorem to Murphy's setting; this is achieved in Scholium 2.9.

Extending Murphy's result beyond the totally ordered case is problematic, in part because for an arbitrary isometric representation V , the algebra $C^*(V)$ generated by $\{V_s : s \in P\}$ need not be spanned by monomials of the form $V_r V_s^*$: one cannot simplify a product $V_r V_s^* V_t V_u^*$ unless s and t are comparable. In [13], Nica made the important observation that for a large class of partially ordered groups (G, P) , which he called *quasi-lattice ordered* groups, this problem does not exist when V is the left regular representation of P on $\ell^2(P)$: the range projections $V_s V_s^*$ respect the lattice structure of P in the sense that $V_s V_s^* V_t V_t^* = V_{s \vee t} V_{s \vee t}^*$, and this additional algebraic relation allows the simplification $V_r V_s^* V_t V_u^* = V_{rs^{-1}(s \vee t)} V_{ut^{-1}(s \vee t)}^*$. Nica called isometric representations with this property *covariant*, and defined $C^*(G, P)$ as the C^* -algebra which is universal for covariant isometric representations.

Since covariance is automatic when (G, P) is a total order, the class of algebras $C^*(G, P)$ includes those considered by Murphy; in [11], Laca and Raeburn used semigroup crossed product techniques to characterize the faithful representations of $C^*(G, P)$, thus generalizing Murphy's theorem. Motivated by this work, Fowler and Raeburn defined a notion of covariance for representations of a product system E over a quasi-lattice order P , defined a universal algebra $C_{\text{cov}}^*(P, E)$, and characterized its faithful representations when E has finite-dimensional fibers [9].

For systems with infinite-dimensional fibers the definition of covariance involves infinite sums which converge σ -weakly but not in norm; consequently, as observed in [9, Remark 4.5], one cannot be certain that every representation of $C_{\text{cov}}^*(P, E)$ is the integrated form of a covariant representation. In Section 1 we give a concrete illustration of this pathology (Example 1.3), and then define a class of product systems, called *compactly aligned*, for which there is no such problem: there is a C^* -algebraic characterization of covariance (Proposition 1.4), and this assures that $C_{\text{cov}}^*(P, E)$ behaves as a universal object should (Proposition 1.7).

Most of the known examples of discrete product systems are compactly aligned; indeed, if E has finite-dimensional fibers, or if (G, P) is a total order, then E is compactly aligned, and the class of compactly-aligned product systems is closed under free products (Proposition 1.10). One can also

adapt Dihn's construction of von Neumann discrete product systems to an arbitrary (G, P) , and these product systems are compactly aligned as well (Proposition 1.12).

By showing that compactly-aligned product systems satisfy the spanning condition required in [9, Theorem 6.1], we obtain a sufficient condition for faithfulness of a covariant representation (Theorem 2.1), and in Corollary 2.2 we use this condition to generalize a simplicity result of Laca and Raeburn. To obtain a condition which is also necessary, we then specialize to product systems over free products of totally-ordered amenable groups; for such a product system E our main result, Theorem 2.3, characterizes the faithful representations of $C_{\text{cov}}^*(P, E)$. The proof is a modification of the one given in [9, Theorem 5.1]: the gauge action gives an expectation on $C_{\text{cov}}^*(P, E)$, and the key steps are to implement this expectation spatially, and to prove faithfulness on the fixed-point algebra. The details are necessarily different; in particular, in performing the usual norm estimates we need the free product structure to implement Cuntz's technique of using an aperiodic sequence to kill the off-diagonal terms.

We would like to thank Iain Raeburn for many helpful discussions during the preparation of this work.

1. COMPACTLY-ALIGNED DISCRETE PRODUCT SYSTEMS

We begin by recalling the pertinent definitions from [9]. Suppose P is a countable discrete semigroup with identity e . A *product system* over P is a family $p : E \rightarrow P$ of nontrivial separable complex Hilbert spaces $E_t := p^{-1}(t)$ which is endowed with an associative multiplication $E \times E \rightarrow E$ in such a way that p is a semigroup homomorphism, and such that for every $s, t \in P$ the map $x \otimes y \in E_s \otimes E_t \mapsto xy \in E_{st}$ extends to a unitary isomorphism. We also insist that $\dim E_e = 1$, so that E has an identity Ω [9, Lemma 1.3].

A *representation* of E in a unital C^* -algebra B is a map $\phi : E \rightarrow B$ which satisfies

- (i) $\phi(xy) = \phi(x)\phi(y)$ for every $x, y \in E$; and
- (ii) $\phi(y)^*\phi(x) = \langle x, y \rangle 1$ if $p(x) = p(y)$.

It follows that the restriction of ϕ to each E_t is linear and isometric, and that $\phi(\Omega) = 1$ [9, Remarks 1.8]. Condition (ii) implies that each $\phi(x)$ is a multiple of an isometry; moreover, the image of an orthonormal basis for E_t is a Toeplitz-Cuntz family, and this collection of families indexed by P generates $C^*(\phi(E))$. When $B = \mathcal{B}(\mathcal{H})$ for some separable Hilbert space $\mathcal{H}$, we call ϕ a *representation of E on $\mathcal{H}$* .

Given a representation $\phi : E \rightarrow B$, for each $s \in P$ there is a homomorphism $\rho_s^\phi : \mathcal{K}(E_s) \rightarrow B$ which satisfies

$$\rho_s^\phi(x \otimes \overline{y}) = \phi(x)\phi(y)^*, \quad x, y \in E_s;$$

here $x \otimes \overline{y}$ denotes the rank-one operator $z \mapsto \langle z, y \rangle x$. It is easy to check that

$$(1.1) \quad \rho_s^\phi(S)\phi(z) = \phi(Sz) \quad \text{for } S \in \mathcal{K}(E_s), z \in E_s,$$

and if $\sigma : B \rightarrow C$ is a homomorphism, then

$$(1.2) \quad \rho_s^{\sigma \circ \phi}(S) = \sigma(\rho_s^\phi(S)) \quad \text{for } S \in \mathcal{K}(E_s).$$

When $B = \mathcal{B}(\mathcal{H})$, ρ_s^ϕ has a unique extension to $\mathcal{B}(E_s)$; we write ρ_s^ϕ for this extension as well, and remark that it is normal (σ -weakly continuous). By [9, Proposition 1.11], there is also an associated semigroup homomorphism $t \mapsto \alpha_t^\phi$ of P into the $*$ -endomorphisms of $\mathcal{B}(\mathcal{H})$, given by

$$(1.3) \quad \alpha_t^\phi(A) = \sum_{u \in \mathcal{B}} \phi(u) A \phi(u)^* \quad \text{for } A \in \mathcal{B}(\mathcal{H}),$$

where the sum converges strongly over any orthonormal basis $\mathcal{B}$ for E_t .

The following Lemma summarizes the properties of α^ϕ and ρ^ϕ we shall require.

Lemma 1.1. *If $s, t \in P$, $z \in E_s$, $S \in \mathcal{B}(E_s)$ and $A \in \mathcal{B}(\mathcal{H})$, then*

- (1) $\alpha_t^\phi(1) = \rho_t^\phi(1)$;
- (2) $\phi(z) \alpha_t^\phi(A) = \alpha_{st}^\phi(A) \phi(z)$;
- (3) $\rho_s^\phi(S) \phi(z) = \phi(Sz)$;
- (4) $\rho_{st}^\phi(S \otimes 1) = \rho_{st}^\phi(1) \rho_s^\phi(S) = \rho_s^\phi(S) \rho_{st}^\phi(1)$; and
- (5) $\rho_{ts}^\phi(1 \otimes S) = \alpha_t^\phi(\rho_s^\phi(S))$.

Remark 1.2. We have written $S \otimes 1$ for the operator satisfying $S \otimes 1(xy) = (Sx)y$ for $x \in E_s$ and $y \in E_t$. If more precision is needed we write $S \otimes 1^t$. Similarly, $1 \otimes S(yx) = y(Sx)$.

Proof of Lemma 1.1. (1) If $\mathcal{B}$ is an orthonormal basis for E_t , then both $\alpha_t^\phi(1)$ and $\rho_t^\phi(1)$ equal $\sum_{u \in \mathcal{B}} \phi(u) \phi(u)^*$.

(2) See [9, Lemma 3.6].

(3) Take a bounded net $\{K_\lambda\}$ of compact operators which converges strongly to S . Then $K_\lambda \rightarrow S$ σ -weakly as well, so (3) follows from (1.1).

(4) By linearity and σ -weak continuity we may assume that $S = x \otimes \bar{y}$. The series $\sum_{u \in \mathcal{B}} xu \otimes \bar{y}u$ converges σ -weakly to $S \otimes 1$, so

$$\begin{aligned} \rho_{st}^\phi(S \otimes 1) &= \sum_{u \in \mathcal{B}} \phi(xu) \phi(yu)^* = \phi(x) \left(\sum_{u \in \mathcal{B}} \phi(u) \phi(u)^* \right) \phi(y)^* \\ &= \phi(x) \alpha_t^\phi(1) \phi(y)^* = \alpha_{st}^\phi(1) \phi(x) \phi(y)^* = \rho_{st}^\phi(1) \rho_s^\phi(S). \end{aligned}$$

The proof that $\rho_{st}^\phi(S \otimes 1) = \rho_s^\phi(S) \rho_{st}^\phi(1)$ is similar.

(5) Again we may assume that $S = x \otimes \bar{y}$, and

$$\alpha_t^\phi(\rho_s^\phi(S)) = \sum_{u \in \mathcal{B}} \phi(u) \phi(x) \phi(y)^* \phi(u)^* = \rho_{ts}^\phi \left(\sum_{u \in \mathcal{B}} ux \otimes \bar{y}u \right) = \rho_{ts}^\phi(1 \otimes S).$$

□

We are primarily interested in the case where P is embedded in a group G , and (G, P) is *quasi-lattice ordered* in the sense of Nica [13]; see also [11, §1] and [9, §3]. Briefly, this means that $P \cap P^{-1} = \{e\}$, so that $s \leq t$ iff $s^{-1}t \in P$ defines a (left-invariant) partial order on G , and that in this partial order every finite subset of G which has an upper bound in P has a least upper bound in P . When $s, t \in P$ have a common upper bound we write $s \vee t$ for the least upper bound; when s and t have no common upper bound we write $s \vee t = \infty$. Our main examples will be direct sums and free products of totally-ordered amenable groups.

If (G, P) is a quasi-lattice ordered group and E is a product system over P , we say that a representation ϕ of E on $\mathcal{H}$ is *covariant* if

$$(1.4) \quad \rho_s^\phi(1)\rho_t^\phi(1) = \begin{cases} \rho_{s \vee t}^\phi(1) & \text{if } s \vee t < \infty \\ 0 & \text{otherwise.} \end{cases}$$

(By Lemma 1.1(1), this definition agrees with [9, Definition 3.2].) By [9, Theorem 4.3], there is a pair $(C_{\text{cov}}^*(P, E), i_E)$ consisting of a unital C^* -algebra $C_{\text{cov}}^*(P, E)$ and a representation $i_E : E \rightarrow C_{\text{cov}}^*(P, E)$ which is universal for covariant representations of E , in the sense that

(a) there is a faithful unital representation σ of $C_{\text{cov}}^*(P, E)$ on Hilbert space such that $\sigma \circ i_E$ is a covariant representation of E ;

(b) for every covariant representation ϕ of E there is a unital representation ϕ_* of $C_{\text{cov}}^*(P, E)$, called the *integrated form* of ϕ , such that $\phi = \phi_* \circ i_E$; and

(c) $C_{\text{cov}}^*(P, E)$ is generated as a C^* -algebra by $i_E(E)$.

Condition (a) falls short of assuring that every representation of $C_{\text{cov}}^*(P, E)$ is the integrated form of a covariant representation. The following example shows that there are product systems for which this deficiency is unavoidable.

Example 1.3. Let $\mathcal{E}$ be a Hilbert space with basis $\{e_k : k \in \mathbb{N}\}$, and let E be the product system over $\mathbb{N} \oplus \mathbb{N}$ whose fiber over (m, n) is $\{(m, n)\} \times \mathcal{E}^{\otimes(m+n)} \cong \mathcal{E}^{\otimes(m+n)}$, and with multiplication given by

$$((k, l), \xi)((m, n), \eta) := ((k + m, l + n), \xi \otimes \eta).$$

If $\{S_k : k \in \mathbb{N}\} \subset \mathcal{B}(\mathcal{H})$ is a collection of isometries with mutually orthogonal ranges, there is a representation ϕ^S of E which satisfies

$$\phi^S((m, n), e_{k_1} \otimes e_{k_2} \otimes \cdots \otimes e_{k_{m+n}}) := S_{k_1} S_{k_2} \cdots S_{k_{m+n}}.$$

The projection $L_{m,n}$ onto the closed linear span of $\phi^S(E_{m,n})\mathcal{H}$ depends only on $m + n$. Moreover, the function $m + n \mapsto L_{m,n}$ is strictly decreasing if $\sum S_k S_k^* < 1$, and is constant if $\sum S_k S_k^* = 1$. Hence ϕ^S is covariant if and only if $\sum S_k S_k^* = 1$.

Suppose then that $\sum S_k S_k^* = 1$. Since ϕ^S is covariant, there is a representation $\phi_*^S : C_{\text{cov}}^*(P, E) \rightarrow \mathcal{B}(\mathcal{H})$ such that $\phi_*^S \circ i_E = \phi^S$. Let $\{T_k : k \in \mathbb{N}\}$ be another Toeplitz-Cuntz family with $\sum T_k T_k^* < 1$. By Cuntz's uniqueness theorem for $\mathcal{O}_\infty$, there is a representation σ of $\phi_*^S(C_{\text{cov}}^*(P, E)) = C^*(S_k)$ such that $\sigma(S_k) = T_k$. By the previous paragraph, $(\sigma \circ \phi_*^S) \circ i_E = \sigma \circ \phi^S = \phi^T$ is not covariant, and hence $\sigma \circ \phi_*^S$ is not the integrated form of a covariant representation.

The following Proposition hints at how to avoid this kind of pathology.

Proposition 1.4. *Suppose (G, P) is a quasi-lattice ordered group and E is a product system over P . A representation $\phi : E \rightarrow \mathcal{B}(\mathcal{H})$ is covariant if and only if, whenever $s, t \in P$, $S \in \mathcal{K}(E_s)$, and $T \in \mathcal{K}(E_t)$, we have*

$$(1.5) \quad \rho_s^\phi(S)\rho_t^\phi(T) = \begin{cases} \rho_{s \vee t}^\phi((S \otimes 1)(T \otimes 1)) & \text{if } s \vee t < \infty \\ 0 & \text{if } s \vee t = \infty. \end{cases}$$

Proof. If ϕ is covariant, then

$$\begin{aligned}\rho_s^\phi(S)\rho_t^\phi(T) &= \rho_s^\phi(S)\rho_s^\phi(1)\rho_t^\phi(1)\rho_t^\phi(T) \\ &= \rho_s^\phi(S)\rho_{s\vee t}^\phi(1)\rho_t^\phi(T) = \rho_{s\vee t}^\phi((S \otimes 1)(T \otimes 1)),\end{aligned}$$

where the last equality uses Lemma 1.1(4). The converse follows by σ -weak continuity. $\square$

If the product $(S \otimes 1)(T \otimes 1)$ in (1.5) were known to be compact, this characterization of covariance would make sense when ϕ maps into an abstract C^* -algebra. This motivates the following Definition.

Definition 1.5. Suppose (G, P) is a quasi-lattice ordered group and E is a product system over P . We say that E is *compactly aligned* if whenever $s, t \in P$ have a common upper bound and S and T are compact operators on E_s and E_t , respectively, the operator $(S \otimes 1)(T \otimes 1)$ is a compact operator on $E_{s\vee t}$. If E is compactly aligned and ϕ is a representation of E in a C^* -algebra B , we say that ϕ is *covariant* if (1.5) holds whenever $s, t \in P$, $S \in \mathcal{K}(E_s)$ and $T \in \mathcal{K}(E_t)$.

Remark 1.6. If E has finite-dimensional fibers, or if (G, P) is a total order, then E is obviously compactly aligned.

The following Proposition shows that the pathology observed in Example 1.3 cannot occur when E is compactly aligned; every representation of $C_{\text{cov}}^*(P, E)$ is the integrated form of a covariant representation.

Proposition 1.7. *Let E be a compactly-aligned product system over P .*

- (1) *The universal map $i_E : E \rightarrow C_{\text{cov}}^*(P, E)$ is covariant.*
- (2) *If σ is a representation of $C_{\text{cov}}^*(P, E)$, then $\sigma \circ i_E$ is covariant.*

Proof. (1) Let σ be a faithful representation of $C_{\text{cov}}^*(P, E)$ on a Hilbert space such that $\phi := \sigma \circ i_E$ is covariant. Since E is compactly aligned we can apply (1.2) to both sides of (1.5) to obtain

$$(1.6) \quad \sigma(\rho_s^{i_E}(S)\rho_t^{i_E}(T)) = \begin{cases} \sigma(\rho_{s\vee t}^{i_E}((S \otimes 1)(T \otimes 1))) & \text{if } s \vee t < \infty \\ 0 & \text{if } s \vee t = \infty. \end{cases}$$

The result follows by applying σ^{-1} to this equation.

(2) Taking $\phi = i_E$ in (1.5) and applying σ to both sides gives (1.6); covariance of $\sigma \circ i_E$ then follows from (1.2). $\square$

For compactly-aligned product systems, we can improve on [9, Proposition 3.7]; the series given there converges in *norm*:

Proposition 1.8. *Let E be a compactly-aligned product system over P , and suppose $v, w \in E$ satisfy $p(v) \vee p(w) < \infty$. Let $\mathcal{B}$ and $\mathcal{C}$ be orthonormal bases for $E_{p(v)^{-1}(p(v) \vee p(w))}$ and $E_{p(w)^{-1}(p(v) \vee p(w))}$, respectively.*

- (1) *The series*

$$\sum_{f \in \mathcal{B}, g \in \mathcal{C}} \langle wg, vf \rangle vf \otimes \overline{wg}$$

converges in norm to $((v \otimes \overline{v}) \otimes 1)((w \otimes \overline{w}) \otimes 1) \in \mathcal{K}(E_{p(v) \vee p(w)})$.

(2) If $\phi : E \rightarrow B$ is covariant, then the series

$$\sum_{f \in \mathcal{B}, g \in \mathcal{C}} \langle wg, vf \rangle \phi(f) \phi(g)^*$$

converges in norm to $\phi(v)^* \phi(w)$.

Proof. Since E is compactly aligned, $K := ((v \otimes \bar{v}) \otimes 1)((w \otimes \bar{w}) \otimes 1)$ is compact. The finite sums $\sum 1 \otimes (f \otimes \bar{f})$ are finite-rank projections which increase to 1, so the series $\sum_{f \in \mathcal{B}} (1 \otimes (f \otimes \bar{f}))K$ converges in norm to K . Similarly, the series $\sum_{g \in \mathcal{C}} K(1 \otimes (g \otimes \bar{g}))$ converges in norm to K . Hence

$$\sum_{f \in \mathcal{B}, g \in \mathcal{C}} (1 \otimes (f \otimes \bar{f}))K(1 \otimes (g \otimes \bar{g}))$$

converges in norm to K . Since

$$\begin{aligned} (1 \otimes (f \otimes \bar{f}))K(1 \otimes (g \otimes \bar{g})) &= ((v \otimes \bar{v}) \otimes (f \otimes \bar{f}))((w \otimes \bar{w}) \otimes (g \otimes \bar{g})) \\ &= (vf \otimes \bar{v}\bar{f})(wg \otimes \bar{w}\bar{g}) \\ &= \langle wg, vf \rangle vf \otimes \bar{w}\bar{g}, \end{aligned}$$

this gives (1).

(2) The series

$$\sum_{f \in \mathcal{B}, g \in \mathcal{C}} \langle wg, vf \rangle \phi(vf) \phi(wg)^* = \sum_{f \in \mathcal{B}, g \in \mathcal{C}} \langle wg, vf \rangle \rho_{p(v) \vee p(w)}^\phi(vf \otimes \bar{w}\bar{g})$$

converges in norm to $\rho_{p(v) \vee p(w)}^\phi(K)$, which by the covariance of ϕ is equal to

$$\rho_{p(v)}^\phi(v \otimes \bar{v}) \rho_{p(w)}^\phi(w \otimes \bar{w}) = \phi(v) \phi(v)^* \phi(w) \phi(w)^*.$$

Multiplying on the left by $\phi(v)^*$ and on the right by $\phi(w)$ gives (2). $\square$

The analysis of many C^* -algebras is facilitated by the existence of a set of Wick-ordered monomials which have dense linear span. The following Corollary shows that when E is compactly aligned, $C_{\text{cov}}^*(P, E)$ is such an algebra.

Corollary 1.9. *If E is a compactly-aligned product system over P , then*

$$C_{\text{cov}}^*(P, E) = \overline{\text{span}}\{i_E(x)i_E(y)^* : x, y \in E\}.$$

Proof. Taking $\phi = i_E$ in Proposition 1.8(2) shows that the right hand side is closed under multiplication. The result follows since each generator $i_E(x)$ of $C_{\text{cov}}^*(P, E)$ can be written as $i_E(x) = i_E(x)i_E(\Omega)^*$, where Ω is the identity of E . $\square$

Since our results in the next section involve free products, we need the following Proposition. (See [9, Examples 1.4(E4)] for the definition of the free product.)

Proposition 1.10. *For each λ belonging to some index set Λ , let (G^λ, P^λ) be a quasi-lattice ordered group, and let E^λ be a product system over P^λ . If each E^λ is compactly aligned, then so is their free product $*E^\lambda$.*

Proof. Suppose $s, t \in *P^\lambda$ have a common upper bound, $S \in \mathcal{K}(E_s)$, and $T \in \mathcal{K}(E_t)$. If $s \leq t$, then the product $(S \otimes 1)(T \otimes 1)$ reduces to $(S \otimes 1)T$, and is hence compact; the case $t \leq s$ is similar. If $s, t < s \vee t$, then there exists $\mu \in \Lambda$ such that $s^{-1}t \in G^\mu$; that is, we can write $s = rs'$, $t = rt'$ with $r \in *P^\lambda$, $s', t' \in P^\mu$ and $s' \vee t' < \infty$. By linearity and continuity we may assume that $S = R_1 \otimes S'$, where $R_1 \in \mathcal{K}(E_r)$ and $S' \in \mathcal{K}(E_{s'})$; similarly, we assume that $T = R_2 \otimes T'$. Then

$$\begin{aligned} (S \otimes 1)(T \otimes 1) &= ((R_1 \otimes S') \otimes 1)((R_2 \otimes T') \otimes 1) \\ &= R_1 R_2 \otimes ((S' \otimes 1)(T' \otimes 1)) \end{aligned}$$

is compact since E^μ is compactly aligned. $\square$

Remark 1.11. In [9, Examples 1.4], it was also shown how one can twist a product system E by a multiplier $\omega : P \times P \rightarrow \mathbb{T}$; it is easy to see that E^ω is compactly aligned if E is. One can also take tensor products of product systems, and this also preserves the property of being compactly aligned.

We close this section by showing how to adapt Dinh's construction of von Neumann discrete product systems [6, Example 6.1] to an arbitrary left-cancellative semigroup P . Fix a separable Hilbert space $\mathcal{H}$ and a unit vector $\xi \in \mathcal{H}$. For $s \in P$, let $\mathcal{H}_s$ be the von Neumann tensor product $\otimes_{P \setminus sP} \mathcal{H}$ with canonical unit vector ξ ; that is, $\mathcal{H}_s$ is the Hilbert space inductive limit $\varinjlim \otimes_F \mathcal{H}$, taken over the finite subsets $F \subseteq P \setminus sP$, under the isometric embeddings obtained by tensoring with ξ [14]. Vectors of the form $\otimes_{r \in P \setminus sP} \eta_r$, where $\eta_r = \xi$ for all but finitely many r , have dense linear span in $\mathcal{H}_s$.

Let $E := \bigsqcup_{s \in P} \{s\} \times \mathcal{H}_s$, $p(s, \eta) := s$, and define multiplication in E by

$$\left(s, \bigotimes_{r \in P \setminus sP} \eta_r\right) \left(t, \bigotimes_{r \in P \setminus tP} \zeta_r\right) := \left(st, \bigotimes_{r \in P \setminus stP} \gamma_r\right),$$

where

$$\gamma_r := \begin{cases} \eta_r & \text{if } r \in P \setminus sP, \\ \zeta_a & \text{if } r = sa \text{ for some } a \in P \setminus tP. \end{cases}$$

It is routine to check that $p : E \rightarrow P$ is a product system. Let d be the dimension of $\mathcal{H}$. Since any other pair $(\mathcal{H}', \xi')$ with $\dim \mathcal{H}' = d$ determines a product system which is isomorphic to the one above, we denote this product system E^d , and call it a *von Neumann discrete product system*.

Proposition 1.12. *If (G, P) is a quasi-lattice ordered group, then any von Neumann discrete product system over P is compactly aligned.*

Proof. Suppose $S \in \mathcal{K}(E_s)$, $T \in \mathcal{K}(E_t)$, and $s \vee t < \infty$; we will show that $(S \otimes 1)(T \otimes 1)$ is compact. Since $P \setminus sP = (P \setminus (sP \cup tP)) \sqcup (tP \setminus sP)$, there is a canonical isomorphism

$$E_s \cong \left(\bigotimes_{P \setminus (sP \cup tP)} \mathcal{H} \right) \otimes \left(\bigotimes_{tP \setminus sP} \mathcal{H} \right).$$

Similarly, there is a canonical isomorphism

$$E_t \cong \left(\bigotimes_{P \setminus (sP \cup tP)} \mathcal{H} \right) \otimes \left(\bigotimes_{sP \setminus tP} \mathcal{H} \right).$$

By linearity and continuity, we may assume that under these identifications $S = S_1 \otimes S_2$ and $T = T_1 \otimes T_2$, where S_1, S_2, T_1 , and T_2 are compact. Using the decomposition $P \setminus (s \vee t)P = (P \setminus (sP \cup tP)) \sqcup (tP \setminus sP) \sqcup (sP \setminus tP)$, we find that $S \otimes 1 = S_1 \otimes S_2 \otimes 1$ and $T \otimes 1 = T_1 \otimes 1 \otimes T_2$, so

$$(S \otimes 1)(T \otimes 1) = S_1 T_1 \otimes S_2 \otimes T_2$$

is compact. $\square$

2. FAITHFUL REPRESENTATIONS

Proposition 1.8 allows us to apply the main results of [9] to compactly-aligned product systems. For this we need a technical amenability hypothesis on (G, P) ; namely, we require the existence of a homomorphism $\theta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ of quasi-lattice ordered groups such that $\mathcal{G}$ is amenable and, whenever $s \vee t < \infty$, we have

$$(2.1) \quad \theta(s \vee t) = \theta(s) \vee \theta(t) \quad \text{and} \quad \theta(s) = \theta(t) \Rightarrow s = t.$$

This hypothesis is satisfied trivially if G is an amenable group, and if (G, P) is a free product of quasi-lattice ordered groups (G^λ, P^λ) with each G^λ amenable, we can take θ to be the canonical map $*(G^\lambda, P^\lambda) \rightarrow \oplus(G^\lambda, P^\lambda)$ [11, Proposition 4.3].

Theorem 2.1. *Suppose (G, P) is a quasi-lattice ordered group which admits a homomorphism $\theta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ as above. If E is a compactly-aligned product system over P and ϕ is a covariant representation of E on Hilbert space such that*

$$(2.2) \quad \prod_{k=1}^n (1 - \rho_{s_k}^\phi(1)) \neq 0 \quad \text{whenever } s_1, \dots, s_n \in P \setminus \{e\},$$

then the integrated form ϕ_* is faithful on $C_{\text{cov}}^*(P, E)$.

Proof. Since E is compactly aligned, Proposition 1.8 implies that the hypotheses of [9, Theorem 6.1] are satisfied. This in turn implies that the hypotheses of [9, Theorem 5.1] are satisfied, so (2.2) implies that $\pi_\phi \times \phi$ is a faithful representation of $B_P \rtimes_{\tau, E} P$. Since ϕ_* is the restriction of $\pi_\phi \times \phi$ to the subalgebra $C_{\text{cov}}^*(P, E)$ (see [9, Theorem 4.3]), it is faithful when (2.2) holds. $\square$

The following Corollary generalizes [11, Corollaries 5.2 and 5.3]. The second part involves quasi-lattice orders (G, P) which satisfy the following condition:

$$(2.3) \quad \begin{aligned} &\text{for each finite subset } F \subseteq P \setminus \{e\}, \\ &\text{there exists } s \in P \setminus \{e\} \text{ such that } s < s \vee t \text{ for every } t \in F. \end{aligned}$$

For example, one could take G to be a countable dense subgroup of $\mathbb{R}$ with positive cone $P = G \cap [0, \infty)$; one could also take a direct sum of such (G, P) .

Corollary 2.2. *For each λ belonging to some index set Λ , let (G^λ, P^λ) be a quasi-lattice ordered group with G^λ amenable, and let E^λ be a compactly-aligned product system over P^λ . Then $C_{\text{cov}}^*(\ast P^\lambda, \ast E^\lambda)$ is simple if*

- (1) Λ is infinite, or if
- (2) $|\Lambda| \geq 2$, and there exists $\mu \in \Lambda$ such that (G^μ, P^μ) satisfies (2.3).

Proof. By Proposition 1.10, $\ast E^\lambda$ is compactly aligned, so by Proposition 1.7, every representation of $C_{\text{cov}}^*(\ast P^\lambda, \ast E^\lambda)$ is the integrated form of a covariant representation. Since the canonical homomorphism $\theta : \ast(G^\lambda, P^\lambda) \rightarrow \oplus(G^\lambda, P^\lambda)$ satisfies (2.1), Theorem 2.1 applies, and it suffices to verify that (2.2) holds for every covariant representation ϕ .

Suppose $s_1, \dots, s_n \in \ast P^\lambda \setminus \{e\}$. Express $s_k = r_k s'_k$ with $s'_k \in \ast P^\lambda$ and $r_k \in P^{\lambda_k} \setminus \{e\}$ for some $\lambda_k \in \Lambda$. Since $\rho_{s_k}^\phi(1) \leq \rho_{r_k}^\phi(1)$, it suffices to show that

$$(2.4) \quad \prod_{k=1}^n (1 - \rho_{r_k}^\phi(1)) \neq 0.$$

Suppose (1) holds. Then there exists $\lambda \in \Lambda \setminus \{\lambda_k\}$, and for any $t \in P^\lambda \setminus \{e\}$ we have $t \vee r_k = \infty$ for each k . Since ϕ is covariant we deduce that $\rho_t^\phi(1) \prod_{k=1}^n (1 - \rho_{r_k}^\phi(1)) = \rho_t^\phi(1) \neq 0$, and this implies (2.4).

Next suppose (2) holds. By reordering if necessary, assume that there exists $m \leq n$ such that $r_k \in P^\mu$ if and only if $1 \leq k \leq m$. By (2.3), there exists $r \in P^\mu \setminus \{e\}$ such that $r < r \vee r_k$ for $1 \leq k \leq m$; observe that $r \vee r_k = \infty$ if $k > m$. By the covariance of ϕ and Lemma 1.1,

$$\rho_r^\phi(1) \prod_{k=1}^n (1 - \rho_{r_k}^\phi(1)) = \alpha_r^\phi \left(\prod_{k=1}^m (1 - \rho_{r^{-1}(r \vee r_k)}^\phi(1)) \right).$$

Now $|\Lambda| \geq 2$, so there exists $\lambda \in \Lambda \setminus \{\mu\}$. Let $t \in P^\lambda \setminus \{e\}$. Then $t \vee s = \infty$ for every $s \in P^\mu \setminus \{e\}$, so multiplying both sides of the previous equation by $\alpha_r^\phi(\rho_t^\phi(1))$ gives

$$\begin{aligned} \alpha_r^\phi(\rho_t^\phi(1)) \prod_{k=1}^n (1 - \rho_{r_k}^\phi(1)) &= \alpha_r^\phi \left(\rho_t^\phi(1) \prod_{k=1}^m (1 - \rho_{r^{-1}(r \vee r_k)}^\phi(1)) \right) \\ &= \alpha_r^\phi(\rho_t^\phi(1)) \neq 0. \end{aligned}$$

This implies (2.4), completing the proof. $\square$

When E has infinite-dimensional fibers, (2.2) may not be necessary for faithfulness. For example, if E is the product system over $\mathbb{N}$ such that $\dim E_1 = \infty$, then $C_{\text{cov}}^*(\mathbb{N}, E)$ is the simple C^* -algebra $\mathcal{O}_\infty$, whereas the previous Theorem gives only that a Toeplitz-Cuntz family $\{S_1, S_2, \dots\}$ generates a faithful copy of $C_{\text{cov}}^*(\mathbb{N}, E)$ if $\sum S_k S_k^* < 1$. (See [9, Corollary 1.6 and Examples 5.6(2)] for details.)

To obtain a theorem which completely characterizes the faithful representations of $C_{\text{cov}}^*(P, E)$ when E has infinite-dimensional fibers, we assume that (G, P) is a free product of totally-ordered amenable groups. Then

$$(2.5) \quad s \vee t \in \{s, t, \infty\} \quad \text{for every } s, t \in P,$$

a property we shall use extensively. Notice that (2.5) implies that any product system over P is compactly aligned: if $S \in \mathcal{K}(E_s)$ and $T \in \mathcal{K}(E_t)$ with $s \vee t < \infty$, then the product $(S \otimes 1)(T \otimes 1)$ reduces to either $S(T \otimes 1)$ or $(S \otimes 1)T$, and is hence compact.

Theorem 2.3. *For each λ belonging to some index set Λ , let (G^λ, P^λ) be a totally-ordered amenable group. Let E be a product system over $P := *P^\lambda$, and let ϕ be a covariant representation of E in a unital C^* -algebra B . Then $\phi_* : C_{\text{cov}}^*(P, E) \rightarrow B$ is injective if and only if*

$$(2.6) \quad \prod_{k=1}^n (1 - \rho_{s_k}^\phi(1)) \neq 0 \quad \text{whenever } \{s_1, \dots, s_n\} \text{ is a finite subset}$$

of $P \setminus \{e\}$ and $\dim E_{s_k} < \infty$ for each k .

Remark 2.4. Our proof does not use the full strength of these hypotheses, and so may apply in slightly more generality: we require only that (G, P) satisfies (2.5), and that it admits a homomorphism $\theta : (G, P) \rightarrow (\mathcal{G}, \mathcal{P})$ of the type discussed prior to Theorem 2.1.

One direction of this theorem is easy: (2.6) is satisfied by the left regular representation $l : E \rightarrow \mathcal{B}(\bigoplus_{s \in P} E_s)$ [9, Lemmas 1.10 and 3.5], and since the identity operator on each E_{s_k} is compact, we may apply (1.2) to obtain $1 - \rho_{s_k}^l(1) = l_*(1 - \rho_{s_k}^{i_E}(1))$; thus (2.6) is satisfied by the universal map i_E . Composing with an injective ϕ_* and again using (1.2) gives (2.6).

For the converse, we model our proof after [9, Theorem 5.1], making use of the identification of $C_{\text{cov}}^*(P, E)$ as a subalgebra of $B_P \rtimes_{\tau, E} P$ [9, Theorem 4.3]. Proposition 1.8 and the canonical map $\theta : *(G^\lambda, P^\lambda) \rightarrow \bigoplus (G^\lambda, P^\lambda)$ allow us to conclude from [9, Theorem 6.1] that E is amenable; this means that the canonical conditional expectation Φ_δ on $B_P \rtimes_{\tau, E} P$ is faithful on positive elements [9, p. 189]. Since this expectation satisfies

$$(2.7) \quad \Phi_\delta(i_E(x)i_E(y)^*) = \begin{cases} i_E(x)i_E(y)^* & \text{if } p(x) = p(y) \\ 0 & \text{otherwise,} \end{cases}$$

Corollary 1.9 shows that Φ_δ restricts to a map on $C_{\text{cov}}^*(P, E)$; indeed, (2.7) determines Φ_δ on $C_{\text{cov}}^*(P, E)$.

In Proposition 2.7 we show that ϕ_* is isometric on $\Phi_\delta(C_{\text{cov}}^*(P, E))$, and we construct a spatial version Φ_ϕ of Φ_δ such that $\Phi_\phi \circ \phi_* = \phi_* \circ \Phi_\delta$. A standard argument then shows that ϕ_* is injective: $\phi_*(b) = 0 \Rightarrow \Phi_\phi \circ \phi_*(b^*b) = 0 \Rightarrow \phi_* \circ \Phi_\delta(b^*b) = 0 \Rightarrow \Phi_\delta(b^*b) = 0 \Rightarrow b = 0$.

For the remainder of this paper we will assume that the hypotheses of Theorem 2.3 are satisfied, and that (2.6) holds. We also assume that B is represented faithfully and nondegenerately on a Hilbert space $\mathcal{H}$, and thus regard ϕ as a covariant representation of E on $\mathcal{H}$. This allows us to make use of the endomorphisms α_s^ϕ defined in (1.3), and to extend the domain of ρ_s^ϕ to $\mathcal{B}(E_s)$.

The following two technical lemmas will be needed to prove Proposition 2.7. The first provides a projection which can be used to kill those terms of a finite sum $\sum \phi(x_j)\phi(y_j)^*$ which are “too long”, and the second provides a vector which plays the role of the aperiodic sequence used by Cuntz to kill off-diagonal terms [4].

Lemma 2.5. *Let $a \in P$, and suppose F is a finite subset of E such that $p(x) \not\leq a$ for every $x \in F$.*

(1) *For every $\epsilon > 0$, there is a nonzero projection $Q_\epsilon \in \mathcal{B}(\mathcal{H})$ such that*

$$\|\alpha_a^\phi(Q_\epsilon)\phi(x)\| < \epsilon \quad \text{for every } x \in F.$$

(2) *More specifically, suppose C is a finite subset of $P \setminus \{e\}$ such that $a^{-1}p(x) \in C$ for every $x \in F$ such that $a < p(x)$. Let R_C be the projection $\prod_{r \in C} (1 - \rho_r^\phi(1))$. Then $\alpha_a^\phi(R_C)\phi(x) = 0$ for every $x \in F$, and if $R_C = 0$, then there is a unit vector $y_\epsilon \in E$ such that*

$$\|\alpha_a^\phi(\phi(y_\epsilon)\phi(y_\epsilon)^*)\phi(x)\| < \epsilon \quad \text{for every } x \in F.$$

Proof. We need only prove (2). If $x \in F$ satisfies $a \vee p(x) = \infty$, then for any $Q \in \mathcal{B}(\mathcal{H})$, Lemma 1.1 and the covariance of ϕ give

$$\alpha_a^\phi(Q)\phi(x) = \alpha_a^\phi(Q)\rho_a^\phi(1)\rho_{p(x)}^\phi(1)\phi(x) = 0.$$

Hence we can assume that $a \vee p(x) < \infty$, and thus $a < p(x)$, for every $x \in F$.

If $x \in F$, then $r := a^{-1}p(x) \in C$, and for any $R \in \mathcal{B}(E_r)$ we can use Lemma 1.1 to calculate

$$\begin{aligned} \|\alpha_a^\phi(1 - \rho_r^\phi(R))\phi(x)\| &= \|(\rho_a^\phi(1) - \rho_{ar}^\phi(1^a \otimes R))\rho_{ar}^\phi(1)\phi(x)\| \\ &= \|\rho_{ar}^\phi(1 - (1^a \otimes R))\phi(x)\| \\ (2.8) \quad &= \|\phi((1 - (1^a \otimes R))x)\| \\ &= \|(1^a \otimes (1 - R))x\|; \end{aligned}$$

in particular $\alpha_a^\phi(1 - \rho_r^\phi(1))\phi(x) = 0$. Thus $\alpha_a^\phi(R_C)\phi(x) = 0$ for every $x \in F$.

Suppose $R_C = 0$. Let $C_{\min}$ be the set of minimal elements of C . Since the projections $1 - \rho_r^\phi(1)$ increase in r , we have

$$0 = \prod_{r \in C} (1 - \rho_r^\phi(1)) = \prod_{r \in C_{\min}} (1 - \rho_r^\phi(1)).$$

By (2.6), there exists $r_{\min} \in C_{\min}$ such that $\dim E_{r_{\min}} = \infty$. Let $r_{\max}$ be a maximal element of $\{r \in C : r_{\min} \leq r\}$. If $r \in C$ and $r \not\leq r_{\max}$, then by (2.5) and the maximality of $r_{\max}$ we have $r \vee r_{\max} = \infty$; for any $y \in E_{r_{\max}}$, the covariance of ϕ thus gives

$$\phi(y)^* \rho_r^\phi(1) = \phi(y)^* \rho_{r_{\max}}^\phi(1) \rho_r^\phi(1) = 0.$$

Hence $\phi(y)\phi(y)^* \leq 1 - \rho_r^\phi(1)$, and from (2.8) we deduce that

$$(2.9) \quad \alpha_a^\phi(\phi(y)\phi(y)^*)\phi(x) = 0 \quad \text{if } y \in E_{r_{\max}}, x \in F, \text{ and } a^{-1}p(x) \not\leq r_{\max}.$$

By (2.5), the set $\{r \in C : r \leq r_{\max}\}$ is totally ordered, say $r_{\min} = r_1 < r_2 < \dots < r_{n-1} < r_n = r_{\max}$. For each $k \in \{1, \dots, n\}$, let P_k be a finite-rank projection on E_{r_k} which satisfies

$$\|(1^a \otimes (1 - P_k))x\| < \epsilon \quad \text{for every } x \in F \text{ such that } p(x) = ar_k.$$

By (2.8) and (2.9), it suffices to find a unit vector $y_\epsilon \in E_{r_{\max}}$ such that $\phi(y_\epsilon)\phi(y_\epsilon)^* \leq 1 - \rho_{r_k}^\phi(P_k)$ for every $k \in \{1, \dots, n\}$; that is, such that

$$(2.10) \quad \phi(y_\epsilon)^* \rho_{r_k}^\phi(P_k) = 0 \quad \text{for every } k \in \{1, \dots, n\}.$$

If some $E_{r_k^{-1}r_{k+1}}$ is finite-dimensional, then we can create a new collection of finite-rank projections by removing r_k and replacing P_{k+1} by $P_{k+1} \vee$

$(P_{r_k} \otimes 1^{r_k^{-1}r_{k+1}})$. For any $y \in E_{r_{\max}}$ we have $\phi(y)^* = \phi(y)^* \rho_{r_{k+1}}^\phi(1)$, so Lemma 1.1(4) gives

$$\phi(y)^* \rho_{r_k}^\phi(P_k) = \phi(y)^* \rho_{r_{k+1}}^\phi(1) \rho_{r_k}^\phi(P_k) = \phi(y)^* \rho_{r_{k+1}}^\phi(P_{r_k} \otimes 1^{r_k^{-1}r_{k+1}});$$

hence any y_ϵ which satisfies (2.10) for this new collection also works for the original. We therefore assume that each $E_{r_k^{-1}r_{k+1}}$ is infinite-dimensional.

We will define unit vectors $w_1, \dots, w_n \in E$ recursively such that

$$(2.11) \quad w_k \in E_{r_k}, \quad \text{and} \quad \phi(w_k)^* \rho_{r_j}^\phi(P_{r_j}) = 0 \quad \text{for } j \in \{1, \dots, k\};$$

then $y_\epsilon := w_n$ satisfies (2.10), completing the proof. Since $\dim E_{r_1} = \infty$, we can choose $w_1 \in E_{r_1}$ such that $P_{r_1} w_1 = 0$; then

$$\phi(w_1)^* \rho_{r_1}^\phi(P_{r_1}) = \phi(w_1)^* \rho_{r_1}^\phi((w_1 \otimes \overline{w_1}) P_{r_1}) = 0.$$

Suppose $1 \leq k \leq n-1$ and w_k satisfies (2.11). We claim that there is a unit vector $z \in E_{r_k^{-1}r_{k+1}}$ such that $P_{k+1}(w_k z) = 0$. Given this, $w_{k+1} := w_k z$ satisfies (2.11), since

$$\phi(w_k z)^* \rho_{r_{k+1}}^\phi(P_{r_{k+1}}) = \phi(w_k z)^* \rho_{r_{k+1}}^\phi((w_k z \otimes \overline{w_k z}) P_{r_{k+1}}) = 0,$$

and, for $1 \leq j \leq k$,

$$\phi(w_k z)^* \rho_{r_j}^\phi(P_{r_j}) = \phi(z)^* \phi(w_k)^* \rho_{r_j}^\phi(P_{r_j}) = 0.$$

For the existence of such a z , let $e_1, \dots, e_m$ be an orthonormal basis for the range of P_{k+1} , so that $P_{k+1} = \sum_{l=1}^m e_l \otimes \overline{e_l}$. For each $l \in \{1, \dots, m\}$, let z_l be the unique vector in $E_{r_k^{-1}r_{k+1}}$ such that $((w_k \otimes \overline{w_k}) \otimes 1^{r_k^{-1}r_{k+1}}) e_l = w_k z_l$. Since $E_{r_k^{-1}r_{k+1}}$ is infinite-dimensional, there is a unit vector $z \in E_{r_k^{-1}r_{k+1}}$ such that $\langle z, z_l \rangle = 0$ for each l . Then

$$\begin{aligned} P_{k+1}(w_k z) &= \sum_{l=1}^m (e_l \otimes \overline{e_l}) ((w_k \otimes \overline{w_k}) \otimes 1^{r_k^{-1}r_{k+1}}) (w_k z) \\ &= \sum_{l=1}^m (e_l \otimes \overline{w_k z_l}) (w_k z) = 0. \end{aligned}$$

□

Lemma 2.6. *Let C and D be finite subsets of $P \setminus \{e\}$. If the projection $\prod_{r \in C} (1 - \rho_r^\phi(1))$ vanishes, then for every unit vector $y \in E$ there is a unit vector $z \in E$ such that*

$$(2.12) \quad \phi(yz) \phi(yz)^* \alpha_d^\phi(\phi(yz) \phi(yz)^*) = 0 \quad \text{for every } d \in D.$$

Proof. First suppose D is a singleton $\{d\}$. Since

$$\alpha_d^\phi(\phi(yz) \phi(yz)^*) = \sum_{u \in \mathcal{B}} \phi(uyz) \phi(uyz)^*,$$

where $\mathcal{B}$ is an orthonormal basis for E_d , it suffices to find z such that

$$(2.13) \quad \phi(yz)^* \phi(uyz) = 0 \quad \text{for every } u \in E_d.$$

If $p(y) \vee dp(y) = \infty$, then $\phi(y)^* \phi(uy) = 0$, and (2.13) holds for any vector z . Assume then that $p(y) \vee dp(y) < \infty$, and define $s \in P \setminus \{e\}$ by

$$s := \begin{cases} p(y)^{-1} dp(y) & \text{if } p(y) < dp(y), \\ (dp(y))^{-1} p(y) & \text{if } dp(y) < p(y). \end{cases}$$

Let z_1 be a unit vector in E_s . All we need to achieve (2.13) is unit vector $z_2 \in E$ such that $\phi(z_2)^* \phi(z_1) = 0$; for then we let $z := z_1 z_2$, and calculate

$$\begin{aligned} \phi(yz)^* \phi(uyz) &= \phi(z_2)^* \phi(z_1)^* \phi(y)^* \phi(uy) \phi(z_1) \phi(z_2) \\ &= \begin{cases} \langle uy, yz_1 \rangle \phi(z_2)^* \phi(z_1) \phi(z_2) & \text{if } p(y) < dp(y), \\ \langle uy z_1, y \rangle \phi(z_2)^* \phi(z_1)^* \phi(z_2) & \text{if } dp(y) < p(y) \end{cases} \\ &= 0. \end{aligned}$$

If there exists $t \in P$ such that $s \vee t = \infty$, then any $z_2 \in E_t$ will suffice. On the other hand, if $s \vee t < \infty$ for every $t \in P$, then for every $r \in C$ we have either $r \leq s$ or $s \leq r$, and thus

$$0 = (1 - \rho_s^\phi(1)) \prod_{r \in C} (1 - \rho_r^\phi(1)) = 1 - \rho_t^\phi(1)$$

for some $t \leq s$. The hypothesis (2.6) implies that E_t , and hence E_s , is infinite-dimensional, and taking any $z_2 \in E_s$ orthogonal to z_1 suffices.

Now suppose inductively that the lemma holds for some finite set D , and that $d' \in P \setminus \{e\}$; we will show that the lemma holds for $D \cup \{d'\}$. Fix a unit vector $y \in E$, and let $z \in E$ be a unit vector such that (2.12) holds. By the above, there exists a unit vector $z' \in E$ such that

$$\phi(yzz') \phi(yzz')^* \alpha_{d'}^\phi(\phi(yzz') \phi(yzz')^*) = 0.$$

Since $\phi(yzz') \phi(yzz')^* \leq \phi(yz) \phi(yz)^*$, (2.12) holds with zz' in place of z , and the induction is complete. $\square$

Proposition 2.7. *If ϕ satisfies (2.6), then*

- (1) ϕ_* is isometric on $\Phi_\delta(C_{\text{cov}}^*(P, E))$, and
- (2) there is a contractive linear map Φ_ϕ on $\phi_*(C_{\text{cov}}^*(P, E))$ such that

$$\Phi_\phi \circ \phi_* = \phi_* \circ \Phi_\delta.$$

Proof. (1) By Corollary 1.9 and the continuity of Φ_δ , finite sums of the form

$$X := \sum_j i_E(x_j) i_E(y_j)^*,$$

where $p(x_j) = p(y_j)$ for each j , are dense in $\Phi_\delta(C_{\text{cov}}^*(P, E))$. Hence it suffices to fix such an X and show that $\|\phi_*(X)\| = \|X\|$.

In the proof of [9, Proposition 5.4] it was shown that there exists $a \in P$ such that $\|X\| = \|T_a\|$, where T_a is the operator on E_a defined by

$$T_a := \sum_{\{j: p(x_j) \leq a\}} (x_j \otimes \overline{y_j}) \otimes 1^{p(x_j)^{-1}a}.$$

Let $\epsilon > 0$. By Lemma 2.5(1), there is a nonzero projection $Q_\epsilon \in \mathcal{B}(\mathcal{H})$ such that $\|\alpha_a^\phi(Q_\epsilon)\phi(x_j)\| < \epsilon$ whenever $p(x_j) \not\leq a$. Since $\phi_*(X) = \sum_j \phi(x_j)\phi(y_j)^*$,

$$\begin{aligned} & \left\| \alpha_a^\phi(Q_\epsilon)\phi_*(X) - \alpha_a^\phi(Q_\epsilon) \left(\sum_{\{j:p(x_j) \leq a\}} \phi(x_j)\phi(y_j)^* \right) \right\| \\ & \leq \sum_{\{j:p(x_j) \not\leq a\}} \|\alpha_a^\phi(Q_\epsilon)\phi(x_j)\phi(y_j)^*\| \\ & < \epsilon \left(\sum_{\{j:p(x_j) \not\leq a\}} \|y_j\| \right). \end{aligned}$$

But using Lemma 1.1 we have

$$\begin{aligned} \alpha_a^\phi(Q_\epsilon) \left(\sum_{\{j:p(x_j) \leq a\}} \phi(x_j)\phi(y_j)^* \right) &= \alpha_a^\phi(Q_\epsilon)\rho_a^\phi(1) \left(\sum_{\{j:p(x_j) \leq a\}} \rho_{p(x_j)}^\phi(x_j \otimes \overline{y_j}) \right) \\ &= \alpha_a^\phi(Q_\epsilon) \left(\sum_{\{j:p(x_j) \leq a\}} \rho_a^\phi((x_j \otimes \overline{y_j}) \otimes 1) \right) \\ &= \alpha_a^\phi(Q_\epsilon)\rho_a^\phi(T_a), \end{aligned}$$

and [9, Proposition 1.12(2)] gives $\|\alpha_a^\phi(Q_\epsilon)\rho_a^\phi(T_a)\| = \|T_a\|$, so

$$\|\alpha_a^\phi(Q_\epsilon)\phi_*(X)\| \geq \|T_a\| - \epsilon \left(\sum_{\{j:p(x_j) \not\leq a\}} \|y_j\| \right).$$

Since ϵ was arbitrary it follows that

$$\|\phi_*(X)\| \geq \|\alpha_a^\phi(Q_\epsilon)\phi_*(X)\| \geq \|T_a\| = \|X\|,$$

and hence $\|\phi_*(X)\| = \|X\|$ as required.

(2) By Corollary 1.9, finite sums of the form $X := \sum_j i_E(x_j)i_E(y_j)^*$ are dense in $C_{\text{cov}}^*(P, E)$, so it suffices to fix such an X and prove that

$$(2.14) \quad \|\phi_*(\Phi_\delta(X))\| \leq \|\phi_*(X)\|.$$

It was shown in [9, Proposition 5.5] that there exists $a \in P$ such that $\|\phi_*(\Phi_\delta(X))\| = \|T_a\|$, where

$$T_a := \sum_{\{j:p(x_j)=p(y_j) \leq a\}} (x_j \otimes \overline{y_j}) \otimes 1^{p(x_j)^{-1}a}.$$

We will show that for every $\epsilon > 0$ there is a nonzero projection $Q_\epsilon \in \mathcal{B}(\mathcal{H})$ such that

$$(2.15) \quad \|\alpha_a^\phi(Q_\epsilon)\phi(x_j)\phi(y_j)^*\alpha_a^\phi(Q_\epsilon)\| < \epsilon \quad \text{unless } p(x_j) = p(y_j) \leq a.$$

Given this, let N be the number of j 's such that $p(x_j) = p(y_j) \leq a$ does *not* hold, and proceed as in part (1) to estimate

$$\begin{aligned}
\|\phi_*(X)\| &\geq \|\alpha_a^\phi(Q_\epsilon)\phi_*(X)\alpha_a^\phi(Q_\epsilon)\| \\
&\geq \left\| \alpha_a^\phi(Q_\epsilon) \left(\sum_{\{j: p(x_j)=p(y_j) \leq a\}} \phi(x_j)\phi(y_j)^* \right) \right\| - N\epsilon \\
&= \|\alpha_a^\phi(Q_\epsilon)\rho_a^\phi(T_a)\| - N\epsilon \\
&= \|T_a\| - N\epsilon \\
&= \|\phi_*(\Phi_\delta(X))\| - N\epsilon.
\end{aligned}$$

Letting ϵ decrease to zero gives (2.14).

It therefore suffices to find a projection Q_ϵ satisfying (2.15). For each j such that $p(x_j) \leq a$, $p(y_j) \leq a$, $p(x_j) \neq p(y_j)$ and $p(x_j)^{-1}a \vee p(y_j)^{-1}a < \infty$, define $d_j \in P \setminus \{e\}$ by

$$d_j := \begin{cases} (p(x_j)^{-1}a)^{-1}p(y_j)^{-1}a & \text{if } p(x_j)^{-1}a < p(y_j)^{-1}a, \\ (p(y_j)^{-1}a)^{-1}p(x_j)^{-1}a & \text{if } p(y_j)^{-1}a < p(x_j)^{-1}a. \end{cases}$$

Fix j such that $p(x_j) \leq a$, $p(y_j) \leq a$, and $p(x_j) \neq p(y_j)$. If $Q \in \mathcal{B}(\mathcal{H})$, then by Lemma 1.1(2) we have

$$\alpha_a^\phi(Q)\phi(x_j)\phi(y_j)^*\alpha_a^\phi(Q) = \phi(x_j)\alpha_{p(x_j)^{-1}a}^\phi(Q)\alpha_{p(y_j)^{-1}a}^\phi(Q)\phi(y_j)^*,$$

which vanishes if $p(x_j)^{-1}a \vee p(y_j)^{-1}a = \infty$ since ϕ is covariant. On the other hand, if $p(x_j)^{-1}a \vee p(y_j)^{-1}a < \infty$, then $\alpha_{p(x_j)^{-1}a}^\phi(Q)\alpha_{p(y_j)^{-1}a}^\phi(Q)$ equals either $\alpha_{p(x_j)^{-1}a}^\phi(Q)\alpha_{d_j}^\phi(Q)$ or $\alpha_{p(y_j)^{-1}a}^\phi(Q)\alpha_{d_j}^\phi(Q)$, so $\alpha_a^\phi(Q)\phi(x_j)\phi(y_j)^*\alpha_a^\phi(Q)$ vanishes if $Q\alpha_{d_j}^\phi(Q) = 0$. We conclude that any Q_ϵ which satisfies

$$(2.16) \quad Q_\epsilon\alpha_{d_j}^\phi(Q_\epsilon) = 0 \quad \text{for each } j,$$

and

$$(2.17) \quad \|\alpha_a^\phi(Q_\epsilon)\phi(x_j)\phi(y_j)^*\alpha_a^\phi(Q_\epsilon)\| < \epsilon \quad \text{if } p(x_j) \not\leq a \text{ or } p(y_j) \not\leq a,$$

will also satisfy (2.15).

For the existence of such a projection Q_ϵ , let

$$F := \{x_j : p(x_j) \not\leq a\} \cup \{y_j : p(y_j) \not\leq a\},$$

and let

$$C := \{d_j\} \cup \{a^{-1}p(x_j) : a < p(x_j)\} \cup \{a^{-1}p(y_j) : a < p(y_j)\}.$$

If the projection R_C of Lemma 2.5(2) is nonzero, we take it as Q_ϵ ; the lemma assures that (2.17) holds, and since $R_C \leq 1 - \alpha_{d_j}^\phi(1)$ and $\alpha_{d_j}^\phi(R_C) \leq \alpha_{d_j}^\phi(1)$, (2.16) holds as well. On the other hand, if $R_C = 0$, then for appropriately small $\delta > 0$ the unit vector y_δ provided by Lemma 2.5(2) satisfies

$$(2.18) \quad \|\alpha_a^\phi(\phi(y_\delta)\phi(y_\delta)^*)\phi(x_j)\phi(y_j)^*\alpha_a^\phi(\phi(y_\delta)\phi(y_\delta)^*)\| < \epsilon$$

if $p(x_j) \not\leq a$ or $p(y_j) \not\leq a$. Lemma 2.6 then provides a unit vector z such that

$$\phi(y_\delta z)\phi(y_\delta z)^*\alpha_{d_j}^\phi(\phi(y_\delta z)\phi(y_\delta z)^*) = 0 \quad \text{for each } j,$$

so $Q_\epsilon := \phi(y_\delta z)\phi(y_\delta z)^*$ satisfies (2.16). Since $Q_\epsilon \leq \phi(y_\delta)\phi(y_\delta)^*$, we see from (2.18) that (2.17) holds as well, and the proof is complete. $\square$

Corollary 2.8. *For each λ belonging to some index set Λ , let (G^λ, P^λ) be a totally-ordered amenable group. Let E be a product system over $P := *P^\lambda$. If there exists $\mu \in \Lambda$ such that E_s is infinite-dimensional for every $s \in P^\mu \setminus \{e\}$, then $C_{\text{cov}}^*(P, E)$ is simple.*

Proof. By Proposition 1.7, it suffices to show that every covariant representation ϕ satisfies (2.6). Suppose $s_1, \dots, s_n \in P \setminus \{e\}$ and $\dim E_{s_k} < \infty$ for every k . Take any $s \in P^\mu \setminus \{e\}$, and fix k . Then $s \not\leq s_k$; for otherwise $E_{s_k} \cong E_s \otimes E_{s^{-1}s_k}$, contradicting $\dim E_{s_k} < \infty = \dim E_s$. On the other hand, $\dim E_{s_k} < \infty$ implies $s_k \notin P^\mu \setminus \{e\}$, and hence $s_k \not\leq s$. From (2.5) we conclude that $s \vee s_k = \infty$ for every k . Since ϕ is covariant we thus have $\rho_s^\phi(1) \prod_{k=1}^n (1 - \rho_{s_k}^\phi(1)) = \rho_s^\phi(1) \neq 0$, and this implies (2.6). $\square$

Taking $|\Lambda| = 1$ gives the following generalization of [6, Theorem 2.2].

Scholium 2.9. *Let (G, P) be a totally-ordered amenable group, and let E be a product system over P . If E_s is infinite-dimensional for every $s \in P \setminus \{e\}$, then $C^*(P, E)$ is simple.*

REFERENCES

- [1] S. Adji, M. Laca, M. Nilsen and I. Raeburn, *Crossed products by semigroups of endomorphisms and the Toeplitz algebras of ordered groups*, Proc. Amer. Math. Soc. **122** (1994), 1133–1141.
- [2] W. Arveson, *Continuous analogues of Fock space*, Memoirs Amer. Math. Soc. **80** (1989), No. 409.
- [3] L. A. Coburn, *The C^* -algebra generated by an isometry*, Bull. Amer. Math. Soc. **73** (1967), 722–726.
- [4] J. Cuntz, *Simple C^* -algebras generated by isometries*, Comm. Math. Phys. **57** (1977), 173–185.
- [5] J. Cuntz, *K -theory for certain C^* -algebras*, Ann. Math. **113** (1981), 181–197.
- [6] H. T. Dinh, *Discrete product systems and their C^* -algebras*, J. Funct. Anal. **102** (1991), 1–34.
- [7] H. T. Dinh, *On generalized Cuntz C^* -algebras*, J. Operator Theory **30** (1993), 123–135.
- [8] R. G. Douglas, *On the C^* -algebra of a one-parameter semigroup of isometries*, Acta Math. **128** (1972), 143–152.
- [9] N. J. Fowler and I. Raeburn, *Discrete product systems and twisted crossed products by semigroups*, J. Funct. Anal. **155** (1998), 171–204.
- [10] M. Laca, *Discrete product systems with twisted units*, Bull. Austral. Math. Soc. **52** (1995), 317–326.
- [11] M. Laca and I. Raeburn, *Semigroup crossed products and the Toeplitz algebras of nonabelian groups*, J. Funct. Anal. **139** (1996), 415–440.
- [12] G. J. Murphy, *Ordered groups and Toeplitz algebras*, J. Operator Theory **18** (1987), 303–326.
- [13] A. Nica, *C^* -algebras generated by isometries and Wiener-Hopf operators*, J. Operator Theory **27** (1992), 17–52.
- [14] J. von Neumann, *On infinite direct products*, Comp. Math. **6** (1938), 1–77.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NEWCASTLE, NSW 2308, AUSTRALIA

E-mail address: neal@math.newcastle.edu.au